


James Webb Space Telescope (JWST)



The First Light Machine

JWST Summary

- **Mission Objective**

- Study origin & evolution of galaxies, stars & planetary systems
- Optimized for near infrared wavelength (0.6 – 28 μm)
- 5 year Mission Life (10 year Goal)

- **Organization**

- Mission Lead: Goddard Space Flight Center
- International collaboration with ESA & CSA
- Prime Contractor: Northrop Grumman Space Technology
- Instruments:
 - Near Infrared Camera (NIRCam) – Univ. of Arizona
 - Near Infrared Spectrometer (NIRSpec) – ESA
 - Mid-Infrared Instrument (MIRI) – JPL/ESA
 - Fine Guidance Sensor (FGS) – CSA
- Operations: Space Telescope Science Institute



Today



1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019

Concept Development

Design, Fabrication, Assembly and Test

science operations



Phase A



Phase B



Phase C/D

Launch

Phase E

Formulation
Authorization

ICR
[i.e., PNAR]

T-NAR

PDR/NAR
(Program Commitment)



Formulation

Implementation


...

Origins Theme's Two Fundamental Questions

- How Did We Get Here?
- Are We Alone?



JWST Science Themes



Three Key Facts

There are 3 key facts about JWST that enables it to perform its Science Mission:

It is a Space Telescope



It is an Infrared Telescope

It has a Large Aperture


Why go to Space

Atmospheric Transmission drives the need to go to space.

Infrared (mid and far/sub-mm) Telescopes (also uv, x-ray, and gamma-ray) cannot see through the Atmosphere



Infrared Light



Why Infrared ?



California Nebula in Center of Picture

Pleiades (Seven Sisters) Cluster is close to bottom.

In the visible this
region of space is
black and devoid of
information.

(from Hubble)

In the sub-mm this
region of space is full
of the coldest matter
of the Universe.

(from Planck, ESA)



M-83



Visible light (taken by the Wide Field Imager on the 2.2-metre MPG/ESO telescope at La Silla in Chile) and infrared (as seen by HAWK-I).


In the infrared, the dust that obscures many stars becomes nearly transparent, making the spiral arms less dramatic, but revealing a whole host of new stars that are otherwise invisible.

Image: ESO/M. Gieles.

(Dr. Emily Baldwin, Astronomy Now, 20 May 2010)

Why do we need Large Apertures?

Aperture = Sensitivity




Adapted from *Cosmic Discovery*, M. Harwit


Sensitivity Matters



GOODS CDFS – 13 orbits




HUDF – 400 orbits



JWST will be more Sensitive than Hubble or Spitzer

HUBBLE





2.4-meter
 $T \sim 270 \text{ K}$




$123'' \times 136''$
 $\lambda/D_{1.6\mu\text{m}} \sim 0.14''$

JWST



6.5-meter
 $T \sim 40 \text{ K}$



$132'' \times 164''$
 $\lambda/D_{2\mu\text{m}} \sim 0.06''$



$114'' \times 84''$
 $\lambda/D_{20\mu\text{m}} \sim 0.64''$

SPITZER



0.8-meter
 $T \sim 5.5 \text{ K}$



$312'' \times 312''$
 $\lambda/D_{5.6\mu\text{m}} \sim 2.22''$




$324'' \times 324''$
 $\lambda/D_{24\mu\text{m}} \sim 6.2''$


**JWST 6X more sensitive
with similar resolution**

JWST 44X more sensitive

Wavelength Coverage



How big is JWST?



Full Scale JWST Mockup





21st National Space Symposium, Colorado Springs, The Space Foundation

Full Scale JWST Mockup



21st National Space Symposium, Colorado Springs, The Space Foundation

How JWST Works



JWST Orbits the 2nd Lagrange Point (L2)

239,000 miles (384,000km)



Earth







Moon

930,000 miles (1.5 million km)




L2

JWST Design: Key Features



OTE Architecture Concept



JWST Requirements

Optical Telescope Element


- 25 sq meter Collecting Area
- 2 micrometer Diffraction Limit
- < 50K (~35K) Operating Temp

Primary Mirror

- 6.6 meter diameter (tip to tip)
- < 25 kg/m² Areal Density
- < \$6 M/m² Areal Cost
- 18 Hex Segments in 2 Rings
- Drop Leaf Wing Deployment


Segments

- 1.315 meter Flat to Flat Diameter
- < 20 nm rms Surface Figure Error






Low (0-5 cycles/aper)	4 nm rms
CSF (5-35 cycles/aper)	18 nm rms
Mid (35-65K cycles/aper)	7 nm rms
Micro-roughness	<4 nm rms

Fun Fact – Mirror Surface Tolerance




Technology Development of Large Optical Systems

MSFC is the JWST Primary Mirror Segment Technology Development Lead for JWST



The 18 Primary Mirror segments

JWST Mirror Technology History



Based on lessons learned, JWST invested early in mirror technology to address lower areal densities and cryogenic operations

Advantages of Beryllium

Very High Specific Stiffness – Modulus/Mass Ratio


Saves Mass – Saves Money

High Conductivity & Below 100K, CTE is virtually zero.


Thermal Stability

Figure Change: 30-55K Operational Range

Beryllium




ULE Glass



Mirror Manufacturing Process

Blank Fabrication



Machining of Web Structure

Machining



Machining of Optical Surface




Completed Mirror Blank

Polishing



Mirror System Integration



Brush Wellman




Axsys Technologies


Batch #1 (Pathfinder) PM Segments



PMSA #1 (EDU-A / A1)



PMSA #2 (3 / B1)



PMSA #3 (4 / C1)

Batch #2 PM Segments



PMSA #4 (5 / A2)



PMSA #5 (6 / B2)




PMSA #6 (7 / C2)

Fun Facts – Mirror Manufacturing

Before





After



Over 90% of material is removed to make each mirror segment – want a little mirror with your Be dust?




Mirror Processing at Tinsley




Tinsley Laboratory – Final Shipment






Primary Mirror Segment Assembly at BATC



MSFC Cryogenic Test Facility



XRCF Cryo Test




- Survival Temperature



- Cryo Deployment
- Nominal Measurement
- Hexapod Deformation Pose
- RoC Actuation Test
- Hexapod Envelope Test
- Pullout Current & Redundant Test (3 of 6 PMSAs)

- Set RoC
- Nominal Measurement
- Hexapod Tilt Test
- Pullout Current & Redundant Test (3 of 6 PMSAs)

Flight Mirrors in XRCF







Primary Mirror Cryogenic Tests







Mirror Fabrication Status at L-3 SSG-Tinsley – July 11

ALL DONE & DELIVERED

Spare Mirrors



Flight Mirrors




SURFACE FIGURE ERROR

Total PM Composite: 13.5 nm RMS



Telescope PM Target: 21.2 nm RMS

Gold Coated Mirror Assemblies



Primary Mirror Backplane

Pathfinder backplane (central section) is complete for test procedure verification at JSC
Flight Backplane under construction



OTE Integration

GSFC - JWST Cleanroom



Backplane Support Structure






Robot arm for mirror installation



Watch JWST being built on the web !

<http://jwst.gsfc.nasa.gov/webcam.html>

Passive Cooling: Sunshield Evolution



Sunshield Development

Evolutionary Pathfinder



Bench Test Articles



Deployment tests



**Clearance
check**




Fight Sunshield at NeXolve in Huntsville





Lee Roop, The Huntsville Times, September 20, 2011

Observatory level testing occurs at JSC Chamber A

Verification Test Activities in JSC Chamber-A



Crosscheck Tests in JSC Chamber-A



Chamber A:

- 37m tall, 20m diameter, 12m door
- LN2 shroud and GHe panels

JWST Launched on Ariane 5 Heavy

**JWST folded and stowed for launch
in 5 m dia x 17 m tall fairing**



**Launch from Kourou Launch Center
(French Guiana) to L2**





JWST vs. HST - orbit

NORTHROP GRUMMAN
Space Technology

Sun

Earth

Moon



HST in Low Earth Orbit, ~500 km up.
Imaging affected by proximity to Earth

JWST will operate at the 2nd Lagrange Point (L2) which is 1.5
Million km away from the earth

L2



L2 Orbit Enables Passive Cryogenic Operation




Second Lagrange Point (L2) of Sun-Earth System

This point follows the Earth around the Sun

The orbital period about L2 is ~ 6 months

Station keeping thrusters required to maintain orbit

Propellant sized for 11 years ($\Delta v \sim 93$)




JWST observes whole sky while remaining continuously in shadow of its sunshield

Field of Regard is annulus covering 35% of the sky


Whole sky is covered each year

Deployment Movie

JWST Optical Path



The JWST telescope is a three mirror anastigmat equipped with a fine steering mirror



JWST's is a Three Mirror Anastigmat (TMA)

Optical design: $f/20$

Diameter of PM: 6.6 m

Effective focal length: 131.4 m

Clear aperture area: 25 m²



Field of view: 18.2 x 9.1 arcmin

Elliptical $f/1.2$ Primary Mirror


Hyperbolic Secondary Mirror creates $f/9$ intermediate image

Elliptical Tertiary Mirror images pupil at Fine Steering Mirror


Transmitted Wavefront Error is 131 nm rms



Deployed Telescope Phasing




Processes have been demonstrated on the Test Bed Telescope as part of TRL-6 development



JWST Science Instruments

enable imagery and spectroscopy over the 0.6 – 29 micron spectrum



JWST Science Theme #1

End of the dark ages: first light and reionization

What are the first luminous objects?

What are the first galaxies?

How did black holes form and interact with their host galaxies?


When did re-ionization of the inter-galactic medium occur?

What caused the re-ionization?

... to identify the first luminous sources to form and to determine the ionization history of the early universe.




Hubble Ultra Deep Field

A Brief History of Time



Redshift

The further away an object is, the more its light is **redshifted** from the visible into the infrared.



When and how did reionization occur?

Re-ionization happened at $z > 6$ or
< 1 B yrs after Big Bang.

WMAP says maybe twice?

Probably galaxies, maybe quasar
contribution

Key Enabling Design Requirments:

Deep near-infrared imaging survey
(1nJy)


Near-IR multi-object spectroscopy

Mid-IR photometry and spectroscopy


JWST Observations:

Spectra of the most distant quasars


Spectra of faint galaxies



First Light: Observing Reionization Edge



Lyman Forest Absorption



Patchy Absorption





Black Gunn-Peterson trough

At 780 M yrs after Big Bang the Universe was up to 50% Neutral. But, by 1 B years after BB is was as we see it today.

Neutral 'fog' was dissolved by very bright 1st Generation Stars (5000X younger & ~100X more massive than our sun).

How do we see first light objects?



2.2 Gyr

1.8 Gyr

1.0 Gyr

Oldest Gamma Ray Burst – 520M yrs after BB

29 Apr 2009, SWIFT detected 5 sec gamma ray burst.

Afterglow in Gemini image has no visible light.


Also, no red-shifted Lyman ‘forest’ was detected.

Once afterglow faded, nothing was visible

TOO FAR

Estimated Age is 520 million years after big bang,
13.14 billion light-years from Earth (Red Shift 9.4).

These first light objects are **TOO RED SHIFTED** for
current telescopes. JWST will study them.



Credit: Gemini Observatory



Jonathan Amos Science correspondent, BBC News

Tammy Plotner, Universe Today, May 26, 2011

2nd Generation Stars – 700M yrs after BB

This star is a 2nd generation star after the big bang because it has trace amounts of heavy elements – meaning that at least one supernova had exploded before it was formed.

But its existence contradicts current theories because it has too much Hydrogen and too much Helium and not enough Carbon and other heavy elements.



Nola Taylor Redd, SPACE.com, 31 August 2011; CREDIT: ESO/Digitized Sky Survey 2

When and How did the First Galaxies Form

First galaxies are small & faint


Light is redshifted into infrared.

Low-metallicity, massive stars.


SNe! GRBs!

JWST Observations

Ultra-Deep NIR survey (1.4 nJy),
spectroscopic & Mid-IR
confirmation.




Ultra Deep Field


 ERO $z \sim 1$


 AGN $z = 5.5$

Malhotra *et al.* 2004

 Galaxy $z = 5.8$

 Galaxy $z = 6.7$

 QSO $z = 2.5$

 Galaxy $z = 0.48$


Results from UDF



$Z=0.48$




$Z = 5.5$



$Z = 5.8$




$Z = 6.7$




Images of 21 redshift-6 galaxies taken from the UDF

Hubble Ultra Deep Field – Near Infrared




Near-Infrared image taken with new Wide-Field Camera 3 was acquired over 4 days with a 173,000 second exposure.

Hubble Ultra Deep Field – Near Infrared




47 Galaxies have been observed at 600 to 650 Myrs after BB.

Hubble Ultra Deep Field – Near Infrared




At 480 M yrs after big bang ($z \sim 10$) this is oldest observed galaxy. Discovered using drop-out technique.



Left image is visible light, and the next three in near-infrared filters. The galaxy suddenly pop up in the H filter, at a wavelength of 1.6 microns (a little over twice the wavelength the eye can detect). (Discover, Bad Astronomy, 26 Jan 2011)

Hubble Ultra Deep Field – Near Infrared

Chandra Deep Field South



CREDIT: X-ray: NASA/CXC/U.Hawaii/E.Treister et al;
Optical: NASA/STScI/S.Beckwith et al


Keith Cooper, Astronomy Now, 15 June 2011
Taylor Redd, SPACE.com, 15 June 2011

What came first – Galaxies or Black Holes?

Each of these ancient 700 M yrs after BB galaxies has a black hole.

Only the most energetic x-rays are detected, indicating that the black-holes are inside very young galaxies with lots of gas.

Unexpected “Big Babies”: 800M yrs after BB



Spitzer and Hubble have identified a dozen very old (almost 13 Billion light years away) very massive (up to 10X larger than our Milky Way) galaxies.

At an epoch when the Universe was only ~15% of its present size, and ~7% of its current age.

This is a surprising result unexpected in current galaxy formation models.

Oldest & Brightest Quasar – 770M yrs after BB

This Quasar is 770 million years after Big Bang, is powered by a black hole 2 billion times the mass of our Sun and emits 60 trillion times as much light as the sun. How a black hole became so massive so soon after the Big Bang is unknown.

“It is like finding a 6-foot-tall child in kindergarten,” says astrophysicist Marta Volonteri, at the University of Michigan in Ann Arbor.


The spectra of the light from this (and other early light objects) indicate that the Universe was still filled with significant amounts of neutral hydrogen even 770 Myrs after big bang.



Image of ULAS J1120+0641, a very distant quasar powered by a black hole with a mass 2 billion times that of the sun, was created from images taken from surveys made by both the Sloan Digital Sky Survey and the UKIRT Infrared Deep Sky Survey. The quasar appears as a faint red dot close to the centre.
CREDIT:
ESO/UKIDSS/SDSS

JWST – the First Light Machine

With its 6X larger collecting aperture, JWST will see back in time further than Hubble and explore the Universe's first light.



JWST Science Theme #2:

The assembly of galaxies

How did the heavy elements form?

How is the chemical evolution of the universe related to galaxy evolution?

What powers emission from galaxy nuclei?

When did the Hubble Sequence form?

What role did galaxy collisions play in their evolution?

Can we test hierarchical formation and global scaling relations?

What is relation between Evolution of Galaxies & Growth/Development of Black Holes in their nuclei?

... to determine how galaxies and the dark matter, gas, stars, metals, morphological structures, and active nuclei within them evolved from the epoch of reionization to the present day.

Formation of Heavy Elements

Carl Sagan said that we are all ‘star dust’.

All of the heavy elements which exist in the universe were formed from Hydrogen inside of stars and distributed via supernova explosions. But observations in the visible couldn’t find enough dust.

Dust is cold, therefore, it can only be seen in IR.

Looking in the IR (with Herschel and Spitzer) at Supernova 1987A, 100,000X more dust was seen than in the visible – the total mass of this dust equals about half of our Sun.



Image of Supernova 1987A, taken in the infrared by Herschel and Spitzer, shows some of the warm dust surrounding it.

CREDIT: Pasquale Panuzzo

SPACE.com, Taylor Redd, 7 July 2011

Chemical make-up of Early Universe

1.8 B yr after BB gamma-ray burst illuminates neighboring galaxies yielding spectra of their chemical makeup.

Metals in the early universe are higher than expected – indicating that star formation in the early universe was much higher than current theory.



GRB 090323 was first detected on 23 March 2009 by NASA's Fermi space telescope and then the Swift satellite, shortly followed by the ground-based GROND system (Gamma-Ray burst Optical and Near-infrared Detector) at the MPG/ESO 2.2-metre telescope in Chile, as well as ESO's Very Large Telescope (VLT). The VLT observations revealed that the gamma-ray burst injected light through its host galaxy and another nearby galaxy, which are both seen at a redshift of 3.57, equivalent to 12 billion years ago.


Subaru Deep Field: Ancient Supernova 3.7B yrs after BB

22 of 150 ancient supernovae in 10% of Subaru Deep Field

12 occurred around 3.7B yrs after big bang.

Supernova were 10X more frequent at this time than today.

Supernova helped seed early universe with chemical elements.



Clara Moskowitz, SPACE.com, 05 October 2011


Galaxy Formation – 1.1B yrs after BB

The early universe was smooth and without structure. Clumping began small and grew to form large galaxies. But how and when?

At 1.1 B-yrs after big bang, this might be the youngest observed cluster.

Cluster contains 11 min-galaxies which are all much smaller than the Milky Way. One has a 30 million solar mass black hole.

It's likely that the mini-galaxies fell together and merged to form a galaxy.



Cluster COSMOS-AzTEC3, located in the Sextans, contains 11 mini-galaxies (circled red dots). Cluster is 1.1 billion yrs after Big Bang.

Subaru / NASA / JPL-Caltech

Discovery required observations from: Chandra X-ray, James Clerk Maxwell Sub-MM, Hubble, Subaru, Keck, Spitzer & several Radio Telescopes

(Sky and Telescope, Robert Naeye, 13 Jan 2011)


Dark Matter

Dwarf Galaxy Segue 1 has 1000 small, dim, primordial (ancient) low metallicity stars.

But, based on star motion, it has 3400X more mass than can be observed.

Some stars are moving too fast, the only thing keeping the galaxy together is gravity.

Thus, there is Dark Matter.



Dark Matter Distribution

Current Theory says that to hold galaxies together, Dark Matter should be ‘clumped’ in a central bulge.

But, observations of two dwarf galaxies, Fornax and Sculptor (which are 99% dark matter), show that the dark matter within them is spread out smoothly.




It is possible that dark matter might interact more with ordinary matter than currently thought, allowing the regular matter to stir up the dark matter and spread it out. Alternatively, dark matter might move faster than expected and therefore be less prone to clumping in galactic centers.

Image: ESO/Digitized Sky Survey 2

The Hubble Sequence

Hubble classified nearby (present-day) galaxies into
Spirals and Ellipticals.



The Hubble Space Telescope has
extended this to the distant past.



Where and when did the Hubble Sequence form?

How did the heavy elements form?



Galaxy assembly is a process of hierarchical merging

Components of galaxies have variety of ages & compositions

JWST Observations:

- Wide-area near-infrared imaging survey

- Low and medium resolution spectra of 1000s of galaxies at high redshift


- Targeted observations of galactic nuclei



Distant Galaxies are “Train Wrecks”

Interacting Galaxies

Hubble Space Telescope • ACS/WFC • WFPC2



NASA, ESA, the Hubble Heritage (AURA/STScI)-ESA/Hubble Collaboration, and
A. Evans (University of Virginia, Charlottesville/NRAO/Stony Brook University)

STScI-PRC08-16a

2-736.1 $z = 1.355$

Optical

Infrared



Merging Galaxies = Merging Black Holes

Combined Chandra & Hubble data shows two black holes (one 30M & one 1M solar mass) orbiting each other – separated by 490 light-years. At 160 million light-years, these are the closest super massive black holes to Earth.

Theory says when galaxies collide there should be major disruption and new star formation.

This galaxy has regular spiral shape and the core is mostly old stars.

These two galaxies merged with minor perturbations.

Galaxy NGC3393 includes two active black holes
X-ray: NASA/CXC/SAO/G.Fabbiano et al; Optical: NASA/STScI



Charles Q. Choi, SPACE.com, 31 August 2011

Galaxy Clusters – 2.6B yrs after BB

Galaxy clusters are the largest structures in the universe. Bound together by gravity, they require billions of years to form.

Galaxy Clusters are thought to have started to form around 3 B-yrs after big bang.

At 2.6 B-yrs old, this is not the oldest observed galaxy cluster. But, spectra indicates that stars in its constituent galaxies are 1 B-yrs old. Thus, may have started forming about 1.5 B-yrs after BB.

X-ray data (similar to image) shows glow from cloud of very hot gas that holds cluster together. Again, it takes many years to trap hot gas.



Hubble NIR Image of CL J1449+0856, the most distant mature cluster of galaxies found. Color added from ESO's VLT and NAOJ's Subaru Telescope. CREDIT: NASA, ESA, R. Gobat (SPACE.com 09 March 2011)



JKCS 041 at 3.7 B-yr after BB may be one of the Universe's oldest clusters. In Chandra image, X-ray emission is shown in blue. Image: NASA/CXC/INAF/S.Andreon (Astronomy Now, 10 May 2010)

JWST Science Theme #3:


Birth of stars and protoplanetary systems

How do molecular clouds collapse?

How does environment affect star-formation?

What is the mass distribution of low-mass stars?

What do debris disks reveal about the evolution of terrestrial planets?





... to unravel the birth and early evolution of stars, from infall on to dust-enshrouded protostars, to the genesis of planetary systems.

HARDY

David Hardy

Birth of Stars and Proto-planetary Systems

- What is the role of molecular clouds, cores and their collapse in the evolution of stars and planetary systems?
- How do protostars form and evolve?
- How do massive stars form and interact with their environment?
- How do massive stars impact their environment by halting or triggering further star formation. How do they impact the evolution of disks?
- What is the initial mass function down to planetary masses?
- How do protoplanetary systems form and evolve?
- How do astrochemical tracers track star formation and the evolution of protoplanetary systems?



How do proto-stellar clouds collapse?

Stars form in small regions collapsing gravitationally within larger molecular clouds.


Infrared sees through thick, dusty clouds

Proto-stars begin to shine within the clouds, revealing temperature and density structure.

Key JWST Enabling Requirements:

High angular resolution near- & mid-IR imagery

High angular resolution imaging spectroscopy



Barnard 68 in infrared

How does environment affect star-formation?

Massive stars produce wind & radiation

Either disrupt star formation, or causes it.

Boundary between smallest brown
dwarf stars & planets is unknown


Different processes? Or continuum?

JWST Observations:

Survey dark clouds, “elephant trunks” or
“pillars of creation” star-forming regions



The Eagle Nebula
as seen in the infrared




Spitzer has Found “The Mountains Of Creation”

Michael Werner, “Spitzer Space Telescope”, William H. Pickering Lecture, AIAA Space 2007.


L. Allen, CfA [GTO]

The Mountains Tell Their Tale

Interstellar erosion & star formation propagate through the cloud



**Young (Solar Mass) Stars are
Shown in This Panel**



**Really Young Stars are Shown in
This Panel**

Star Formation in Dust/Gas Cloud



Herschel discovered 700 newly-forming stars condensing along filaments of dust in a never before penetrated dark cloud at the heart of Eagle Nebula. Two areas glowing brightest in icy blue light are regions where large newborn stars are causing hydrogen gas to shine.

Impossible Stars

100 to 150 solar mass stars should not exist but they do.

When a star gets to 8 to 10 solar mass its wind blows away all gas and dust, creating a bubble and stopping its growth (see Herschel Image).

The bubble shock wave is creating a dense 2000 solar mass region in which an ‘impossible’ star is forming. It is already 10 solar mass and in a few 100 thousand years will be a massive 100 to 150 solar mass – making it one of the biggest and brightest in the galaxy.

(Space.com, 6 May 2010)




Image of RCW 120 (ESA),
Discover.com, Ian O'Neill, 7 May 2010

Orion Nebula Protoplanetary Discs



Hubble has discovered 42
protoplanetary discs in the
Orion Nebula

JWST Science Theme #4:

Planetary systems and the origins of life

How do planets form?

How are circumstellar disks like our Solar System?

How are habitable zones established?

... to determine the physical and chemical properties of planetary systems including our own, and to investigate the potential for the origins of life in those systems.

Robert Hurt

Planetary Formation Questions and 2 Models


- ♦ How do planets and brown dwarfs form?
- ♦ How common are giant planets and what is their distribution of orbits?
- ♦ How do giant planets affect the formation of terrestrial planets?
- ♦ What comparisons, direct or indirect, can be made between our Solar System and circumstellar disks (forming solar systems) and remnant disks?
- ♦ What is the source of water and organics for planets in habitable zones?
- ♦ How are systems cleared of small bodies?
- ♦ What are the planetary evolutionary pathways by which habitability is established or lost?
- ♦ Does our solar system harbor evidence for steps on these pathways?

TWO PLANET FORMATION SCENARIOS

Accretion model



Orbiting dust grains accrete into "planetesimals" through nongravitational forces.



Planetesimals grow, moving in near-coplanar orbits, to form "planetary embryos."




Gas-giant planets accrete gas envelopes before disk gas disappears.



Gas-giant planets scatter or accrete remaining planetesimals and embryos.

Gas-collapse model



A protoplanetary disk of gas and dust forms around a young star.



Gravitational disk instabilities form a clump of gas that becomes a self-gravitating planet.



Dust grains coagulate and sediment to the center of the protoplanet, forming a core.




The planet sweeps out a wide gap as it continues to feed on gas in the disk.

History of Known (current) NEO Population

2006

Earth
Crossing

Outside
Earth's
Orbit



Known

- 340,000 minor planets
- ~4500 NEOs
- ~850 Potentially Hazardous Objects (PHOs)

Scott
Manley

Armagh
Observatory

Follow the DUST


Dust disks are durable and omnipresent



The central star of the Helix Nebula, a hot, luminous White Dwarf, shows an infrared excess attributable to a disk in a planetary system which survived the star's chaotic evolution

Planetary System Formation effects Dust


This star has 3 large (10X Jupiter mass) planets (observed by Hubble, Keck & Gemini North) which are causing a huge halo of fine dust particles (indicating lots of colliding objects) around the star. Dust which can be detected by an infrared telescope.



NASA's Spitzer Space Telescope captured this infrared image of a giant halo of very fine dust around the young star HR 8799, located 129 light-years away in the constellation Pegasus. The brightest parts of this dust cloud (yellow-white) likely come from the outer cold disk similar to our own Kuiper belt (beyond Neptune's orbit). The huge extended dust halo is seen as orange-red. Credit: NASA/JPL-Caltech/Univ. of Ariz.

Spiral Arms Hint At The Presence Of Planets

Disk of gas and dust around a sun-like star has spiral-arm-like structures. These features may provide clues to the presence of embedded but as-yet-unseen planets.




Near Infrared image from Subaru Telescope shows disk surrounding SAO 206462, a star located about 456 light-years away in the constellation Lupus. Astronomers estimate that the system is only about 9 million years old. The gas-rich disk spans some 14 billion miles, which is more than twice the size of Pluto's orbit in our own solar system.

Photonics Online 20 Oct 2011

Direct Image of an ExoPlanet being Formed

Image shows the youngest exoplanet yet discovered. Its Star (slightly smaller than our Sun) is only 2 million years old. Dust is accreting (falling) into the new planet leaving a gap in the planetary disk. New planet is $\sim 6X$ mass of Jupiter.





Using the Keck Telescope

Left: The dusty disk around the star LkCa 15. All of the light at this wavelength is emitted by cold dust in the disk; the hole in the center indicates an inner gap.



Right: An expanded view of the central part of the cleared region, showing a composite of two reconstructed images (blue: 2.1 microns; red: 3.7 microns) for LkCa 15 b. The location of the central star is also marked.

Dust in Planetary Systems


Fomalhaut system at 24 μm
(Spitzer Space Telescope)



Simulated JWST image
Fomalhaut at 24 microns



Kalas, Graham & Clampin 2005




Kalas et al 2008

Planets and Dust


Earth has a ‘tail’ of dust particles.

10 to 20 micrometer size particles are slowed or captured by Earth’s gravity and trail behind Earth. The cloud of particles is about 10 million km wide and 40 million km long.



(Wired.com, Lisa Grossman, 8 July 2010)

Radial Velocity Method finds planets close to stars



61 Virginis (61 Vir) has 3 planets inside of Venus's orbit.

From their star, the planets have masses of ~5X, 18X & 24X Earth's mass.

They orbit 61 Virginis in 4, 38 & 124 day periods.

Also, direct Spitzer observations indicate a ring of dust at twice the distance of Neptune from the star.


Bad Astronomy
Orbital schematic credit: Chris Tinney

Direct Imaging detects planets far from their star

HR 8799 has at least 4 planets


3 planets ('c' has Neptune orbit) were first imaged by Hubble in 1998. Image reanalyzed because of a 2007 Keck discovery.

3 outer planets have very long orbits or 100, 200 & 400 years. Multiple detections are required to see this motion.




Kepler Mission

Kepler (launched in 2009) is hunting planets by staring continuously at 165,000 stars looking for dips in their light caused when a planet crosses in front of the star.



One planetary system (Kepler-11) has a star like ours & 6 mini-Neptune size planets



Five of six Kepler-11 exoplanets (all larger than Earth) orbit their star closer than Mercury orbits the sun. One orbits inside Venus.

Credit: NASA/AP (Pete Spotts, Christian Science Monitor.com, 23 May 2011.)



Kepler Planetary Systems

Kepler has discovered over 1200 planets:

800 in single planet systems,

400 in 170 systems with 2 to 6 transiting planets, and

68 potential Earth size planets (54 in Habitable Zone)



Kepler's planet candidates by size.
CREDIT: NASA/Wendy Stenzel
(SPACE.com 2 Feb 2011)

Graphic shows multiple-planet systems as of 2/2/2011. Hot colors to cool colors (red to yellow to green to cyan to blue to gray) indicate big planets to smaller planets. CREDIT: Daniel Fabrycky (SPACE.com, 23 May 2011)

Is There Life Elsewhere in the Galaxy?

Need to multiply these values by $\eta_{\text{Earth}} \times f_B$ to get the number of potentially life-bearing planets detected by a space telescope.

η_{Earth} = fraction of stars with Earth-mass planets in HZ

f_B = fraction of the Earth-mass planets that have detectable biosignatures

Earth-mass planets within these HZ will be *very*

If: $\eta_{\text{Earth}} \times f_B \sim 1$ then $D_{\text{Tel}} \sim 4\text{m}$

$\eta_{\text{Earth}} \times f_B < 1$ then $D_{\text{Tel}} \sim 8\text{m}$


$\eta_{\text{Earth}} \times f_B \ll 1$ then $D_{\text{Tel}} \sim 16\text{m}$

Number of nearby stars capable of hosting

Kepler is finding that η_{Earth} maybe 1.5% to 2.5% (SPACE.COM, 21 Mar 2011)

Thus, an 8-m telescope might find 1 to 3 Earth twins and an 16-m telescope might find 10 to 20 Earth twins.

Number of FGK stars for which SNR=10, R=70 spectrum of Earth-twin could be obtained in <500 ksec



Green bars show the number of FGK stars that could be observed 3x each in a 5-year mission without exceeding 20% of total observing time available to community.

How are habitable zones established?

Source of Earth's H₂O and organics is not known

Comets? Asteroids?


History of clearing the disk of gas and small bodies

Role of giant planets?




JWST Observations:

Comets, Kuiper Belt Objects

Icy moons in outer solar system



Spitzer Spectrum Shows Water Vapor Falling onto Protoplanetary Disk



Michael Werner, "Spitzer Space Telescope", William H. Pickering Lecture, AIAA Space 2007.

Proto-Stars produce Water

In a proto-star 750 light-years away,
Herschel detected:

Spectra of Atomic Hydrogen and Oxygen
are being pulled into the star, and

Water vapor being spewed at 200,000 km
per hour from the poles.

The water vapor freezes and falls back
onto the proto-planetary disk.

Discovery is because Herschel's
infrared sensors can pierce the
dense cloud of gas and dust feeding
the star's formation.



A Protostar and its Polar Jets NASA/Caltech


Other Herschel Data finds enough
water in the outer reaches of the
young star TW Hydrae (175 light-
yrs from Earth) to fill Earth's
oceans several thousand times over.

Mike Wall, SPACE.com; Date: 20 October 2011


Molecular Oxygen discovered in space

Herschel found molecular oxygen in a dense patch of gas and dust adjacent to star-forming regions in the Orion nebula.

The oxygen maybe water ice that coats tiny dust grains.



All of Life's Ingredients Found in Orion Nebula





Herschel Telescope has measured spectra for all the ingredients for life as we know them in the Orion Nebula.

(Methanol is a particularly important molecule)

Search for Habitable Planets

atmosphere



habitability

L. Cook

interior


surface


Search for Life

What is Life?

What does life do?

Life Metabolizes





All Earth life uses chemical energy generated from redox reactions

Life takes advantage of these spontaneous reactions that are kinetically inhibited

Diversity of metabolisms rivals diversity of exoplanets

Bio Markers

Spectroscopic Indicators of Life



Absorption Lines

CO₂

Ozone

Water





“Red” Edge



Example signs of life from chemical spectra.

Credit: NASA JPL

Earth Through Time



Seager

Kasting Sci. Am. 2004
See Kaltenegger et al. 2006
Earth from the Moon

Beyond JWST

Heavy Lift Launch Vehicle enables even larger telescopes
8-m UV/Optical Telescope or
24-m Far-IR Telescope



HST
2.4m aperture
STS/4.7m




JWST
6.5m aperture
Ariane V/5.4m



SAFIR Monolith
8m aperture
Ares V/10m




SAFIR Baseline
10m aperture
Ariane V/5.4m




SAFIR upscale
19.5m aperture
Ares V/10m


R=100 ATLAST Spectrum of 1 Earth-mass Terrestrial Exoplanet at 10 pc



R=500 ATLAST Spectrum of 1 Earth-mass Terrestrial Exoplanet at 10 pc



Detecting Photometric Variability in Exoplanets



Countdown to Launch

JWST is

making excellent technical progress

will be ready for launch ~2017-2019

will be the dominant astronomical
facility for a decade undertaking a
broad range of scientific
investigations


Ariane 5




Learn more at: www.jwst.nasa.gov

New

New



Download for free at:
jwst.gsfc.nasa.gov



Available Now



www.stsci.edu/jwst/science/whitepapers/

1000s of Scientists and Engineers in USA and around the world are working to make JWST.



Any Questions?



